

Nematic liquid crystal phase
in a system of interacting dimers

Ian Jauslin


joint with Elliott H. Lieb

arXiv: 1709.05297

<http://ian.jauslin.org>

Liquid crystals

- Orientational order and positional disorder.




Models

- (Disertori, Giuliani, Jauslin): hard plates in \mathbb{R}^3 .
- [Disertori, Giuliani, 2013]: hard rods in \mathbb{Z}^2 .
- [Ioffe, Velenik, Zahradník, 2006]: hard rods in \mathbb{Z}^2 (variable length).
- [Bricmont, Kuroda, Lebowitz, 1984]: hard needles in \mathbb{R}^2 .
- [Heilmann, Lieb, 1979]: interacting dimers.


Heilmann-Lieb model

[Heilmann, Lieb, 1979]




Heilmann-Lieb model

[Heilmann, Lieb, 1979]



Heilmann-Lieb model

[Heilmann, Lieb, 1979]



Heilmann-Lieb model

- Grand-canonical Gibbs measure:

$$\langle A \rangle_{\mathbf{v}} := \lim_{\Lambda \rightarrow \mathbb{Z}^2} \frac{1}{\Xi_{\Lambda, \mathbf{v}}(z)} \sum_{\underline{\delta} \in \Omega_{\mathbf{v}}(\Lambda)} A(\underline{\delta}) z^{|\underline{\delta}|} \prod_{\delta \neq \delta' \in \underline{\delta}} e^{\frac{1}{2} J \mathbb{1}_{\delta \sim \delta'}}$$

- ▶ Λ : finite box.
- ▶ $\Omega_{\mathbf{v}}(\Lambda)$: non-overlapping dimer configurations satisfying the boundary condition.
- ▶ $z \geq 0$: fugacity.
- ▶ $J \geq 0$: interaction strength.
- ▶ $\mathbb{1}_{\delta \sim \delta'}$ indicator that dimers are adjacent and aligned.

Theorem

For $1 \ll z \ll J$, $\|(x, y)\|_{\text{HL}} := J|x| + e^{-\frac{3}{2}J} z^{-\frac{1}{2}}|y|$,

- Given two vertical edges e_v, f_v , $\langle \mathbf{1}_{e_v} \rangle_v$ is *independent* of e_v and

$$\langle \mathbf{1}_{e_v} \rangle_v = \frac{1}{2}(1 + O(e^{-\frac{1}{2}J} z^{-\frac{1}{2}}))$$

$$\langle \mathbf{1}_{e_v} \mathbf{1}_{f_v} \rangle_v - \langle \mathbf{1}_{e_v} \rangle_v \langle \mathbf{1}_{f_v} \rangle_v = O(e^{-c \text{dist}_{\text{HL}}(e_v, f_v)})$$

- Given two horizontal edges e_h, f_h , $\langle \mathbf{1}_{e_h} \rangle_v$ is *independent* of e_h and

$$\langle \mathbf{1}_{e_h} \rangle_v = O(e^{-3J})$$

$$\langle \mathbf{1}_{e_h} \mathbf{1}_{f_h} \rangle_v - \langle \mathbf{1}_{e_h} \rangle_v \langle \mathbf{1}_{f_h} \rangle_v = O(e^{-3J - c \text{dist}_{\text{HL}}(e_h, f_h)})$$

1D system

- [Heilmann, Lieb, 1979]: mostly vertical dimers.
- *Only* vertical dimers: integrable.
- Given two vertical edges e_v, f_v , $\langle \mathbf{1}_{e_v} \rangle_v$ is *independent* of e_v and

$$\langle \mathbf{1}_{e_v} \rangle_v = \frac{1}{2}(1 + O(e^{-\frac{1}{2}J} z^{-\frac{1}{2}}))$$

$$\langle \mathbf{1}_{e_v} \mathbf{1}_{f_v} \rangle_v - \langle \mathbf{1}_{e_v} \rangle_v \langle \mathbf{1}_{f_v} \rangle_v = O(e^{-c \operatorname{dist}_{1D}(e_v, f_v)})$$

with $\|(x, y)\|_{1D} := e^{-\frac{3}{2}J} z^{-\frac{1}{2}} |y|$.