

Time-evolution of electron emission from a metal surface


Ian Jauslin

joint with **Ovidiu Costin**, **Rodica Costin**, and **Joel L. Lebowitz**

arXiv:1808.00936


<http://ian.jauslin.org>

Field emission




Field emission

$$V(x) = U\Theta(x), \quad E_F = k_F^2 < U$$




Thermal emission

$$V(x) = U\Theta(x), \quad k^2 > U$$




Photonic emission

$$V_t(x) = \Theta(x)(U - E_t x), \quad E_t = 2\epsilon\omega \cos(\omega t)$$




Field emission

$$V(x) = \Theta(x)(U - Ex)$$



Field emission

- [Millikan, Lauritsen, 1928]: experimental plot of the logarithm of the current against $1/E$



Field emission through a triangular barrier

- [Fowler, Nordheim, 1928]: predicted that the current is, for small E ,

$$J \approx CE^2 e^{-\frac{\alpha}{E}}$$

- ([Rokhlenko, 2011]: studied the range of applicability of the approximation, and found more accurate approximations for larger fields.)

Fowler-Nordheim equation

- Schrödinger equation


$$i\partial_t\psi = -\Delta\psi + \Theta(x)(U - Ex)\psi$$

- Fowler-Nordheim: stationary solution: $\psi_{\text{FN}}(x, t) = e^{-ik^2t}\varphi_{\text{FN}}(x)$

$$\varphi_{\text{FN}}(x) = \begin{cases} e^{ikx} + R_E e^{-ikx} & x < 0 \\ T_E \text{Ai}(e^{-\frac{i\pi}{3}}(E^{\frac{1}{3}}x - E^{-\frac{2}{3}}(U - k^2))) & x > 0 \end{cases}$$

R_E and T_E are chosen so that φ_{FN} and $\partial\varphi_{\text{FN}}$ are continuous at $x = 0$.

Fowler-Nordheim equation



Initial value problem

- Initial condition:

$$\psi(x, 0) = \begin{cases} e^{ikx} + R_0 e^{-ikx} & x < 0 \\ T_0 e^{-\sqrt{U-k^2}x} & x > 0 \end{cases}$$

R_0 and T_0 ensure that ψ and $\partial\psi$ are continuous.

- **Theorem:** $\psi(x, t)$ behaves asymptotically like ψ_{FN} :

$$\psi(x, t) = \psi_{\text{FN}}(x, t) + \left(\frac{t}{\tau_E(x)} \right)^{-\frac{3}{2}} + O(t^{-\frac{5}{2}}).$$

Photoemission

- Photoelectric effect: early observations: [Hertz, 1887], [Hallwachs, 1888], [Lenard, 1900].
- When a metal is irradiated with ultra-violet light, electrons are ionized, with kinetic energies at integer multiples of $\hbar\omega$.
- [Einstein, 1905]: interpretation: quanta of light (*photons*) of energy $\hbar\omega$ are absorbed by the electrons, whose kinetic energy is raised by $n\hbar\omega$, and can escape the metal.

Photoemission

- Time dependent potential:

$$V_t(x) = \Theta(x)(U - 2\epsilon\omega \cos(\omega t)x)$$

- Magnetic gauge:

$$\Psi(x, t) := \psi(x, t)e^{-ix\Theta(x)A(t)}, \quad A(t) := \int_0^t ds \, 2\epsilon\omega \cos(\omega s) = 2\epsilon \sin(\omega t)$$

satisfies

$$i\partial_t \Psi(x, t) = ((-i\nabla + \Theta(x)A(t))^2 + \Theta(x)U) \Psi(x, t)$$

Periodic solution

- [Faisal, Kamiński, Saczuk, 2005]

$$\Psi_{\text{FKS}}(x, t) = \begin{cases} e^{ikx} \exp(-ik^2t) + \Psi_R(x, t) & \text{if } x < 0 \\ \Psi_T(x, t) & \text{if } x > 0 \end{cases}$$

$$\Psi_R(x, t) = \sum_{M \in \mathbb{Z}} R_M e^{-iq_M x} \exp(-iq_M^2 t), \quad q_M = \sqrt{k^2 + M\omega}$$

$$\Psi_T(x, t) = \sum_{M \in \mathbb{Z}} T_M e^{ip_M x} \exp\left(-iUt - i \int_0^t d\tau (p_M - A(\tau))^2\right)$$

$$p_M = \sqrt{k^2 - U + M\omega - 2\epsilon^2}$$

- $\Psi(x, t)$, $(-i\nabla + \Theta(x)A(t))\Psi(x, t)$ are continuous.

Initial value problem

- Initial condition:

$$\Psi(x, 0) = \begin{cases} e^{ikx} + R_0 e^{-ikx} & x < 0 \\ T_0 e^{-\sqrt{U-k^2}x} & x > 0 \end{cases}$$

R_0 and T_0 ensure that Ψ and $\partial\Psi$ are continuous.

- **Conjecture** (in progress): $\Psi(x, t)$ behaves asymptotically like Ψ_{FKS} :

$$\psi(x, t) = \psi_{\text{FKS}}(x, t) + \left(\frac{t}{\tau_{\text{FKS}}(x)} \right)^{-\frac{3}{2}} + O(t^{-\frac{5}{2}}).$$

Idea of the proof: field emission

- Laplace transform:

$$\hat{\psi}_p(x) := \int_0^\infty dt e^{-pt} \psi(x, t)$$

- Schrödinger equation:

$$(-\Delta + \Theta(x)V(x) - ip)\psi_p(x) = -i\psi(x, 0), \quad V(x) := U - Ex$$

Solution in Laplace space

- For simplicity, $R_0 \equiv T_0 \equiv 0$.
- Solution:

$$\hat{\psi}_p(x) = \begin{cases} c(p)e^{\sqrt{-ip}x} - \frac{ie^{ikx}}{-ip + k^2} & \text{if } x < 0 \\ d(p)\varphi_p(x) & \text{if } x > 0 \end{cases}$$

with

$$(-\Delta + V(x) - ip)\varphi_p(x) = 0$$

$$\varphi_p(x) = \text{Ai} \left(e^{-\frac{i\pi}{3}} \left(E^{\frac{1}{3}}x - E^{-\frac{2}{3}}(U - ip) \right) \right)$$


Solution in Laplace space

- c and d ensure that $\hat{\psi}_p(x)$ and $\partial\hat{\psi}_p(x)$ are continuous at $x = 0$:

$$c(p) = \frac{i(ik\varphi_p(0) - \partial\varphi_p(0))}{(-ip + k^2)(\sqrt{-ip}\varphi_p(0) - \partial\varphi_p(0))}$$

$$d(p) = -\frac{i}{(\sqrt{-ip} + ik)(\sqrt{-ip}\varphi_p(0) - \partial\varphi_p(0))}.$$

Poles in Laplace plane



Asymptotic behavior

- As $t \rightarrow \infty$:

$$\psi(x, t) = \psi_{\text{FN}}(x, t) + \left(\frac{t}{\tau_E(x)} \right)^{-\frac{3}{2}} + O(t^{-\frac{5}{2}}).$$

- If $k < 0$ (reflected wave), then there is no pole on the imaginary axis, so there is no contribution as $t \rightarrow \infty$.
- Similarly, the transmitted wave in the initial condition does not contribute.

Idea of the proof: photoemission

- In Laplace space:

$$\hat{\Psi}_p(x) := \int_0^\infty dt e^{-pt} \Psi(x, t)$$

the equation is discrete:

$$\mathfrak{f}_n^{(\sigma)}(x) := \hat{\Psi}_{-ik^2 - i\sigma - in\omega}(x), \quad \mathcal{Re}(\sigma) \in \left[-\frac{\omega}{2}, \frac{\omega}{2}\right)$$

$$\begin{aligned} (-\Delta - k^2 - \sigma - n\omega + \Theta(x) (U + 2\epsilon^2)) \mathfrak{f}_n^{(\sigma)}(x) - \Theta(x) 2\epsilon \nabla (\mathfrak{f}_{n+1}^{(\sigma)}(x) - \mathfrak{f}_{n-1}^{(\sigma)}(x)) \\ - \Theta(x) \epsilon^2 (\mathfrak{f}_{n+2}^{(\sigma)}(x) + \mathfrak{f}_{n-2}^{(\sigma)}(x)) = -i\psi(x, 0) \end{aligned}$$

Initial value problem

- This system of ODEs is *integrable* for $x < 0$ and $x > 0$, so we have closed form expressions for a family of solutions $f_n^{(\sigma)}(x)$, parametrized by two sequences $c_n^{(\sigma)}$ and $d_n^{(\sigma)}$:

$$f_n^{(\sigma)}(x) = \begin{cases} c_n^{(\sigma)} e^{-ix\sqrt{k^2+\sigma+n\omega}} + \frac{ie^{ikx}}{\sigma+n\omega} & , x < 0 \\ \frac{\omega}{2\pi} \sum_{m \in \mathbb{Z}} d_m^{(\sigma)} e^{-\kappa_m^{(\sigma)}x} \int_0^{\frac{2\pi}{\omega}} dt e^{-i(n-m)\omega t} e^{\frac{i\epsilon^2}{\omega} \sin(2\omega t) + \kappa_m^{(\sigma)} \frac{4\epsilon}{\omega} \cos(\omega t)} & , x > 0 \end{cases}$$

with

$$\kappa_m^{(\sigma)} := \sqrt{U + 2\epsilon^2 - k^2 - \sigma - m\omega}$$

Initial value problem

- The sequences c_n and d_n are determined by the continuity condition at $x = 0$:

$$\sum_{m \in \mathbb{Z}} G_{n,m}^{(\sigma)} d_m^{(\sigma)} = v_n^{(\sigma)}, \quad c_n^{(\sigma)} = \sum_{m \in \mathbb{Z}} H_{n,m}^{(\sigma)} d_m^{(\sigma)} + w_n^{(\sigma)}.$$

- The long-time behavior of Ψ depends on the singularities of $\hat{\Psi}_p$ with $p \in i\mathbb{R}$.
- Can prove (by solving the equation for $\psi(x, t)$ using a Fourier transform in x) that the Schrödinger equation has a unique solution. This implies that $G^{(\sigma)}$ is invertible for imaginary σ .
- Only singularities on imaginary axis: $-ik^2 + i\omega\mathbb{Z}$.